

Universal properties of dark matter halos and infall model

Igor Tkachev

INR

Brussels, April 30, 2010

- The flux from DM decay

$$F_{DM} = \frac{\Gamma \Omega_{fov}}{8\pi} \int_{los} dr \rho_{DM}(r)$$

- Average DM column density

$$S = \frac{2}{r_*^2} \int_0^{r_*} r' dr' \int dz \rho(\sqrt{r'^2 + z^2})$$

- We can parametrize it as $S \propto \rho_* r_*$
- S looks to be the same, from dwarfs to clusters
- It is essentially insensitive to the choice of fitting DM profile

Boyarsky et. al., 2006

- The flux from DM decay

$$F_{DM} = \frac{\Gamma \Omega_{fov}}{8\pi} \int_{los} dr \rho_{DM}(r)$$

- Average DM column density

$$S = \frac{2}{r_*^2} \int_0^{r_*} r' dr' \int dz \rho(\sqrt{r'^2 + z^2})$$

- We can parametrize it as $S \propto \rho_* r_*$
- S looks to be the same, from dwarfs to clusters
- It is essentially insensitive to the choice of fitting DM profile

Boyarsky et. al., 2006

- The flux from DM decay

$$F_{DM} = \frac{\Gamma \Omega_{fov}}{8\pi} \int_{los} dr \rho_{DM}(r)$$


- Average DM column density

$$S = \frac{2}{r_*^2} \int_0^{r_*} r' dr' \int dz \rho(\sqrt{r'^2 + z^2})$$


- We can parametrize it as $S \propto \rho_* r_*$
- S looks to be the same, from dwarfs to clusters
- It is essentially insensitive to the choice of fitting DM profile

Boyarsky et. al., 2006

ISO vs NFW




Constant DM surface density?




Donato et al., 2009

An evidence in favour of MOND?



Gentile et al., Nature '09


Scaling of DM column density



$$S \propto M_{\text{halo}}^{0.2}$$

Boyarsky et al., 2009

Scaling of DM column density



Napolitano et al., 2010

Infall model

Position of each particle in the halo obey

$$\frac{d^2r}{dt^2} = -\frac{G M(r,t)}{r^2}$$

For initial perturbations with power-law profiles

$$\frac{\delta M_i}{M_i} = \left(\frac{M_0}{M_i} \right)^\epsilon$$


the halo evolves in a self-similar manner, e. g.

$$M(r,t) = M(t)\mathcal{M}(r/R(t))$$

$$r(r_i, t) = R(t)\lambda(t/t_*)$$


$$\rho(r,t) = \frac{M(t)}{R^3(t)} \times F\left(\frac{r}{R(t)}\right)$$

Infall model




Sikivie, Yun Wang & I.T., 1996

Infall model




Sikivie, Yun Wang & I.T., 1996

Infall model




Sikivie, Yun Wang & I.T., 1996

Infall model




Sikivie, Yun Wang & I.T., 1996

Infall model




Sikivie, Yun Wang & I.T., 1996

Infall model




Sikivie, Yun Wang & I.T., 1996

Infall model




Best fit: $R = 1.07 \pm 0.14$ Mpc, $h = 0.71 \pm 0.5$

G. Steigman & I. T., 1998




Diemand and Kuhlen, 2008





Vogelsberger and White, 2010

Infall model



Sikivie, Yun Wang & I.T., 1996



$\rho \propto r^{-2}$ between first inner and outer caustics

$\rho \propto r^{-\gamma}$ inside first inner caustics, where

$$\gamma = \frac{9\epsilon}{3\epsilon + 1}$$

$\gamma \approx 1.1$ for $\epsilon = 0.2$

Sikivie, Yun Wang & I.T., 1996

DM column density in infall model

Qualitative understanding

- Self-similarity implies

$$S \propto \rho_* r_* \propto \rho_{\text{ta}} R_{\text{ta}}$$


- Infall implies

$$R_{\text{ta}} \propto (G t^2 M)^{1/3}$$


- If there are deviations from self-similarity $S \propto c(M) \cdot M^{1/3}$
- Concentration parameter $c = r_*/R_{\text{ta}}$ is a weak function of mass, $c \propto M^{-0.1}$
- Therefore

$$S \propto \frac{M^{0.23}}{t^{4/3}}$$

Concentration parameter



Scaling of DM column density



Infall $\mathcal{S} \propto M_{\text{halo}}^{1/3-0.1}$, best fit $\mathcal{S} \propto M_{\text{halo}}^{0.2}$

Boyarsky et al., 2009

Gravity vs. cosmological expansion

Let mass M creates a gravitational potential $\phi(r)$

- Turn around time

$$t_{\text{ta}} = \frac{1}{\sqrt{2}} \int_0^{R_{\text{ta}}} \frac{dr}{\sqrt{\phi(r) - \phi(R_{\text{ta}})}}$$

- Any gravitational potential of the form

$$\phi(r) = -\frac{GM}{r} F\left(\frac{\rho(r)}{\rho_0}\right)$$

leads to the scaling $S \propto M^{1/3}$

- Example: Schwarzschild-de Sitter potential

$$\phi(r) = -\frac{GM}{r} - \frac{\Lambda r^2}{6} = -\frac{GM}{r} F\left(\frac{\Lambda}{G\rho(r)}\right)$$

Large scale modifications of gravity

- A possible set of consistent (as a spin-2 field theory) large scale modifications of gravity is described by two parameters - scale r_c and a number $0 \leq \alpha \leq 1$ *Dvali et. al., 2007*
- In these models (DGP, "degravitation")

$$\phi_\alpha(r) = -\frac{GM}{r} \pi\left(\frac{r}{r_V}\right)$$


- where the Vainshtein radius:

$$r_V = \left(2GMr_c^{4(1-\alpha)}\right)^{\frac{1}{1+4(1-\alpha)}}$$

- Only for $\alpha = 1/2$

$$\phi(r) = -\frac{GM}{r} F\left(\frac{\rho(r)}{\rho_0}\right)$$

Restrictions on modifications of gravity



Boyarsky & Ruchayskiy, 2010